

Long-Term Follow-Up of a Randomized Controlled Trial Comparing Acceptance and Commitment Therapy and Standard Cognitive Behavior Therapy for Anxiety and Depression

Evan M. Forman
Jena A. Shaw
Elizabeth M. Goetter
James D. Herbert
Jennie A. Park
Drexel University

Erica K. Yuen
Medical University of South Carolina

The present study represents one of the first comparisons of the long-term effectiveness of traditional cognitive behavior therapy (i.e., Beckian cognitive therapy; CT) and acceptance and commitment therapy (ACT). One hundred thirty-two anxious or depressed outpatients were randomly assigned to receive either CT or ACT, and were assessed at posttreatment ($n=90$) and at 1.5-year ($n=91$) follow-up. As previously reported, the two treatments were equivalently effective at posttreatment according to measures of depression, anxiety, overall (social/occupational/symptom-related) functioning, and quality of life. However, current results suggest that treatment gains were better maintained at follow-up in the CT condition. Clinical significance analyses revealed that, at follow-up, one-third more CT patients were in the clinically normative range in terms of depressive symptoms and more than twice as many CT patients were in the normative range in

terms of functioning levels. The possible long-term advantage of CT relative to ACT in this population is discussed.

Keywords: acceptance-based behavior therapy; psychotherapy outcome; depression; anxiety; long-term follow-up

THE TERM COGNITIVE BEHAVIORAL therapy (CBT) reflects a broad collection of evidence-based approaches that have become the most widely utilized and researched of all psychotherapeutic methods (Norcross, Hedges, & Castle, 2002), with Beckian cognitive therapy (CT; Beck, 1991) representing the most widely used and empirically supported form of CBT (Butler, Chapman, Forman, & Beck, 2006; Hofmann & Smits, 2008). A newer subcategory of CBT, sometimes referred to as acceptance-based behavior therapies, has risen to prominence in recent years. Examples include mindfulness-based cognitive therapy (MBCT; Z. V. Segal, Williams, & Teasdale, 2002), mindfulness-based stress reduction (Kabat-Zinn, 1990), acceptance-based behavior therapy for generalized anxiety disorder (Roemer & Orsillo, 2005), dialectical behavior therapy (DBT; Linehan, 1993), and acceptance and commitment therapy (ACT; Hayes,

Address correspondence to Evan Forman, Ph.D., Department of Psychology, Drexel University, 245 N 15th Street, MS 626, Philadelphia, PA 19102; e-mail: evan.forman@drexel.edu.

0005-7894/43/801–811/\$1.00/0

© 2012 Association for Behavioral and Cognitive Therapies. Published by Elsevier Ltd. All rights reserved.


Strosahl, & Wilson, 1999), among others. Of these, ACT has received the most attention in terms of empirical study (Hayes, Levin, Plumb, Boulanger, & Pistorello, in press; Hayes, Luoma, Bond, Masuda, & Lillis, 2006) and scientific debate (Arch & Craske, 2008; Corrigan, 2001, 2002; Gaudiano, 2009a, 2009b; Hayes, 2002, 2008; Hayes et al., in press; Herbert & Forman, in press, 2011; Hofmann & Asmundson, 2008; Hofmann & Asmundson, in press; Öst, 2008, 2009). At the level of technology, there are some important differences in how ACT and CT treat psychopathology (Forman & Herbert, 2009). CT makes use of cognitive disputation and other forms of reappraisal (including behavioral experiments) designed to correct systematic biases in information processing, with the goal of reducing symptom intensity (Beck, Rush, Shaw, & Emery, 1979). The goal of ACT is *not* symptom reduction per se, but helping patients to “accept” difficult internal experiences (thoughts, images, emotions, sensations) in the service of engaging in values-consistent behavior change.

ACT has demonstrated preliminary effectiveness across a range of problem behaviors, including mood (Zettle & Hayes, 1986) and anxiety (Block, 2003; L. A. Brown et al., 2011; Dalrymple & Herbert, 2007; Roemer, Salters-Pedneault, & Orsillo, 2006; Twohig, Hayes, & Masuda, 2006) disorders, psychosis (Bach & Hayes, 2002; Gaudiano & Herbert, 2006), polysubstance abuse (Hayes, Wilson, et al., 2004), and smoking (R. A. Brown, Lejuez, Kahler, Strong, & Zvolensky, 2005; Hernández-López, Luciano Soriano, Bricker, Roales-Nieto, & Montesinos Marin, 2009; Hernandez Lopez, Roales Nieto, Luciano Soriano, & Montesinos Marin, 2005), among others. A meta-analysis (Hayes, et al., 2006) reported ACT to be superior to active treatments, including standard CBT. However, Öst (2008) has criticized the rigors of the trials on which the meta-analysis relied, and a subsequent meta-analysis (Powers, Zum Vörde Sive Vörding, & Emmelkamp, 2009) concluded that ACT was equally effective as established treatments (but see also Gaudiano, 2009a). The RCT upon which the current study was based similarly detected no differences in efficacy between ACT and CT at posttreatment in the treatment of depression and anxiety (Forman, Herbert, Moitra, Yeomans, & Geller, 2007).

It is important to examine longer-term effectiveness of psychotherapies, as immediate effects can fade over time, patients might take time to master skills learned in treatment, and treatments that are equivalent at one time point can diverge later (Gifford et al., 2004; Lappalainen et al., 2007). Generally speaking, standard CBT has demonstrated longer-term efficacy (Butler et al., 2006; Gloaguen, Cottraux, Cucherat, &

Blackburn, 1998; Shapiro et al., 1994), though several recent reviews suggest effects weaken considerably after 1 year (e.g., Durham, Higgins, Chambers, Swan, & Dow, 2011). Open trials and trials comparing ACT to nonactive treatments or treatment-as-usual support ACT's lasting benefits for anxiety (Dalrymple & Herbert, 2007; Ossman, Wilson, Storaasli, & McNeill, 2006; Twohig, 2008; Zettle, 2003), depression (Blackledge & Hayes, 2006), trichotillomania (Woods, Wetterneck, & Flessner, 2006), psychosis (Bach & Hayes, 2002; Gaudiano & Herbert, 2006), substance dependence (Hayes, Wilson, et al., 2004; Stotts, Masuda, & Wilson, 2009; Twohig, Shoenberger, & Hayes, 2007), smoking (Gifford et al., 2004), obesity (Forman, Butryn, Hoffman, & Herbert, 2009; Lillis, Hayes, Bunting, & Masuda, 2009; Tapper et al., 2009), and chronic pain (Dahl, Wilson, & Nilsson, 2004; McCracken, MacKichan, & Eccleston, 2007; Vowles & McCracken, 2008). However, the follow-up periods of these studies tended to be relatively short (i.e., 1 to 3 months), limiting the extent to which conclusions can be drawn about longer-term effects. Moreover, few of these trials compared ACT to another well-established, active intervention, and none compared ACT to traditional CBT.

The current study compares long-term (18-month follow-up) outcomes from a group of outpatients randomly assigned to receive either CT or ACT. The study is an extension of an earlier report of posttreatment outcomes (Forman, Herbert, et al., 2007). As mentioned, the effects of standard CBT may attenuate in the longer-term (Durham et al., 2011), and some proponents have hypothesized that ACT might hold certain advantages over standard (i.e., Beckian) CBT because ACT is argued to be more tightly linked to basic research on its mechanisms and underlying theory (Hayes, 2008; Hayes et al., in press). ACT proponents have also questioned the putative mechanisms of change of CT (i.e., modification of the content of dysfunctional cognitions; Hayes, 2008; Hayes, Villatte, Levin, & Hildebrandt, 2011). On the other hand, standard CBT has proven long-term efficacy (Butler et al., 2006; Gloaguen et al., 1998; Shapiro et al., 1994). Perhaps an overriding consideration is the accumulating evidence suggesting that it is the behavioral elements of treatment that represent the mechanisms of action, and that other components are superfluous (Dimidjian et al., 2006; Longmore & Worrell, 2007). Thus, no specific hypotheses were made regarding differential long-term effectiveness of the two treatments. Given previous findings that mindfulness and acceptance variables moderate the impact of treatment, we tentatively hypothesized such a moderation effect at follow-up.



† No significant differences in age, gender, marital status, ethnicity, or condition were found between participants with or without follow-up data

FIGURE 1 Consort diagram. *Note.* CT = Cognitive Therapy; ACT = Acceptance and Commitment Therapy.

Method

PARTICIPANTS

Participants were 132 nonbaccalaureate health science students presenting for treatment at a student counseling center (Figure 1). Inclusion criteria were purposefully broad to maximize external validity. Individuals were eligible if they were seeking individual psychotherapy (as opposed to, for example, couples therapy, or study skills training), reported at least moderate levels of anxiety and/or depression (i.e., score of 9 or above on the Beck Depression Inventory [BDI-II; Beck, Steer, & Brown, 1996] or Beck Anxiety Inventory [BAI; Beck, Epstein, Brown, & Steer, 1988]),¹ were fluent in English, and were not psychotic.

Participants ranged in age from 18 to 52 ($M = 26.7$, $SD = 6.4$). The majority were female (79.5%) and white (70.7%); half (50.7%) were single, and a small

number were taking psychiatric medication (16.7%).² Diagnoses were obtained by a structured interview, the Mini-International Neuropsychiatric Interview (MINI), which is a widely used diagnostic interview with shorter administration times, but comparable reliability and validity (Lecrubier et al., 1997; Sheehan et al., 1998), to the Structured Clinical Interview for DSM Disorders (SCID; First, Williams, Spitzer, & Gibbon, 2007). The large majority of patients met criteria for anxiety disorders (49.2%; 24.2% generalized anxiety disorder, 12.1% anxiety disorder NOS, 5.3% panic disorder, 5.3% social anxiety disorder, 3.1% OCD, 2.3% specific phobia, 1.5% PTSD) and/or depressive disorders (37.1%; 22% recurrent MDD, 9.8% depressive disorder NOS, 7.6% single episode MDD, 3.0% dysthymia). Other diagnoses included adjustment disorders (6.8%) and eating disorders (3.8%). A

¹ We elected to use a cutoff score of ≥ 9 on the BDI-II and BAI to indicate elevated levels of depression and anxiety, respectively, based on data examining normative data for adult samples on the BDI-II (D. L. Segal, Coolidge, Cahill, & O'Riley, 2008; Whisman, Perez, & Ramel, 2000) and BAI (Gillis, Haaga, & Ford, 1995).

² Seven participants who reported taking psychiatric medications at baseline reported that they no longer were taking medications at follow-up. As a check, the analyses described below were repeated without these 7 participants and results were equivalent.

diagnostic category (depressive disorder, anxiety disorder, both depressive and anxiety disorders or neither depressive nor anxiety disorder) was established for analytical purposes. Diagnostic assessors had received 4 hours of training on the MINI, along with a minimum of 3 hours of supervised practice. Additionally, diagnostic interviews were recorded and were reviewed by doctoral-level supervisors.

PROCEDURE

After providing consent to participate in the study, patients were assigned to either the ACT ($n=69$) or CT ($n=63$) condition via stratified block randomization determined by total score on the Outcome Questionnaire (OQ; Lambert, Hansen, et al., 1996). As the course of therapy was intended to be naturalistic, there was not a standardized number of sessions for either condition; termination was based on mutual consensus between the therapist and patient. However, the average number of sessions across conditions was similar for ACT (18.10, $SD=15.91$) and CT (16.37, $SD=13.64$; $t(130)$, $p=.50$). Both interventions included psychoeducation, coping skills development, and behavioral (especially behavioral exposure) exercises that were presented within the theoretical framework of the assigned intervention. CT incorporated discussion of automatic thoughts, core beliefs, and schemas, identification of cognitive distortions, cognitive disputation, and cognitive restructuring, whereas ACT emphasized experiential acceptance, mindfulness training, clarification of personal values, and willingness to experience internal distress for the sake of living consistently with one's values.

Data were collected at baseline, posttreatment, and at approximately 18-month³ follow-up. Individuals were paid \$20.00 for completing the follow-up procedures, but not for any of the other assessments.

Therapist Allegiance, Fidelity, and Competence

Therapists ($n=28$) were doctoral-level graduate students who received specialized training (a 30-hour workshop) and weekly supervision in both ACT and CT. The training and supervision sessions were led by licensed clinical psychologists with several years' combined experience in ACT and CT. Therapists administered treatments to both ACT and CT patients. As might be expected given the dual training model, no significant differences in therapist allegiance were detected in terms of which treatment therapists

judged to be more effective (ACT=53%; $\chi^2=1.29$, $p=.26$) or preferred (CT=61%; $\chi^2=.14$, $p=.71$). Session-by-session treatment manuals were not utilized; instead, therapists followed general ACT or CT treatment outlines that were specific to diagnosis. All sessions were audio recorded and a random selection of 3 sessions per study participant were rated using the Drexel University ACT and CT Adherence and Rating Scale (McGrath, Forman, & Herbert, 2009), revealing that adherence, contamination, and competence were statistically equivalent between groups. Specifically, an average of 36.2% (ACT) and 33.3% (CT) of time was spent on treatment-specific components, with the majority of the remaining time spent on common treatment components, e.g., behavioral interventions and nonspecific, active listening/supportive interventions. Contamination rates were low: 4.3% (ACT) and 5.7% (CT) of time was spent on aspects associated with the nonassigned treatment condition. Competence of therapist was rated as "good," "very good," or "excellent" in 91% of ACT sessions and 93% of CT sessions.

MEASURES

Beck Depression Inventory – 2nd Edition (BDI-II; Beck et al., 1996)

The BDI-II is an extensively used 21-item assessment tool, designed to assess the severity of current depressive symptoms. The BDI-II has been shown to have good reliability and strong content, concurrent, and discriminant validity in both clinical and nonclinical samples (Beck, Steer, & Garbin, 1988).

Beck Anxiety Inventory (BAI; Beck, Epstein, et al., 1988)

The BAI assesses the severity of anxiety related symptoms. The BAI is a 21-item self-report measure with high internal consistency ($\alpha=.92$) and good reliability and validity in clinical populations (Beck et al.).

Outcome Questionnaire (OQ; Lambert, Hansen, et al., 1996)

The OQ is designed to measure patient functioning in subjective distress, interpersonal relationships, and social role performance. The 45 items yield a total score from 0 to 180, with higher scores indicating poorer functioning. The OQ has demonstrated excellent internal consistency and appropriate content and concurrent validity (Lambert, Burlingame, et al., 1996).

Quality of Life Index (QOLI; Frisch et al., 2005)

The QOLI is a measure of life satisfaction rooted in the view that overall life quality is the sum of satisfaction in a variety of life domains. Clients are asked to rate the importance of (0=*not important*

³ Follow-up assessments varied from 14 to 20 months (mode=18 months). The nature of the sample was such that most participants had left the region, and locating them proved to be slow and arduous, producing variability in follow-up times. Due to this variability, time to follow-up was entered as a covariate in all analyses.

Table 1
Baseline Differences in Demographic Variables

		Completed follow-up (%)		No follow-up (%)		Comparisons using logistic regression (Wald's χ^2 (p))		
		ACT	CT	ACT	CT	Follow-up	Condition	Follow x Condition
Gender	Female	82.6	80.0	69.6	83.3	0.55 (.46)	1.01 (.31)	1.02 (.31)
Marital Status	Married	47.8	55.6	39.1	61.1	0.42 (.52)	1.92 (.17)	0.57 (.45)
Ethnicity	White	76.1	68.9	73.9	50.0	1.32 (.25)	2.42 (.12)	0.69 (.41)
Comorbidity	Present	30.4	40.0	21.7	32.3	0.01 (.92)	0.68 (.41)	0.04 (.84)

Note. CT = Cognitive Therapy; ACT = Acceptance and Commitment Therapy.

to 2 = *extremely important*) and satisfaction with (-3 = *very dissatisfied* to 3 = *very satisfied*) a variety of life domains. The index takes the sum of the products of each domain. Test-retest coefficients and internal consistency coefficients are very good (Frisch et al.).

Kentucky Inventory of Mindfulness Skills (KIMS; Baer, Smith, & Allen, 2004)

The KIMS is a 39-item measure of four components of mindfulness: observing, describing, acting with awareness, and accepting without judgment (Baer et al., 2004). Items are rated on a 5-point Likert scale ranging from 1 (*never or very rarely true*) to 5 (*almost always or always true*). The measure was found to have high internal consistency, adequate to good test-retest reliability, and validation analyses providing support for the relationship between mindfulness and mental health (Baer et al.).

Acceptance and Action Questionnaire (AAQ; Hayes, Strosahl, et al., 2004)

The AAQ is a 9-item measure of the extent to which an individual demonstrates an accepting attitude towards negative feelings and experiences and the

ability to take action even when feeling dysphoric or uncertain. Items are rated on a 1 (*never true*) to 7 (*always true*) scale, with higher scores indicating greater levels of experiential avoidance. The AAQ has demonstrated very good internal consistency, and has adequate criterion-related, predictive, and convergent validities (Hayes et al.).

Global Assessment of Functioning (GAF) and Clinical Global Impression (CGI)

At baseline and posttest, but not at follow-up, clinicians rated GAF (Spitzer, Gibbon, Williams, & Endicott, 1996), which ranges from 1 (*persistent danger of hurting oneself or others*) to 100 (*superior functioning*), and also the CGI (Guy, 1976), which ranges from 1 (*normal*) to 7 (*extreme illness*).

Results

PARTICIPANT ENROLLMENT

Key variables, including primary diagnostic category, $\chi^2(1) = .69, p = .88$, and days to follow-up ($t = -.20, p = .84$), were equivalent by assignment to condition. In addition, there were no significant differences between those who provided follow-up data ($n = 91$)

Table 2
Baseline Differences by Follow-up Status

	Completed follow-up (M (SD))		No follow-up (M (SD))		Comparisons using ANOVA (F (p))		
	ACT	CT	ACT	CT	Follow-up	Condition	Interaction
BDI	19.2 (10.4)	19.2 (10.2)	16.5 (7.7)	17.2 (5.8)	1.80 (.18)	0.04 (.85)	0.03 (.85)
BAI	15.0 (10.4)	15.5 (11.2)	14.3 (13.6)	10.5 (8.7)	1.88 (.17)	0.63 (.42)	1.03 (.31)
OQ	81.4 (16.7)	82.3 (16.8)	75.1 (15.3)	82.3 (13.5)	1.07 (.30)	1.74 (.19)	1.05 (.31)
QOLI	1.04 (2.49)	0.51 (2.08)	0.80 (1.71)	0.25 (2.10)	0.34 (.56)	1.66 (.20)	<0.01 (.99)
AAQ	53.0 (5.6)	53.5 (6.8)	53.0 (7.9)	54.2 (8.9)	0.07 (.79)	0.43 (.51)	0.07 (.80)
KIMS-Observe	25.0 (8.5)	24.9 (9.7)	22.4 (7.6)	22.4 (6.8)	2.46 (.12)	<0.01 (.97)	<0.01 (.99)
KIMS-Describe	18.6 (6.6)	20.1 (7.0)	19.4 (5.9)	16.7 (6.0)	1.04 (.31)	0.24 (.63)	2.77 (.10)
KIMS- Act with Awareness	17.6 (6.3)	17.6 (6.6)	17.8 (4.1)	16.4 (5.0)	0.19 (.66)	0.41 (.53)	0.36 (.55)
KIMS- Acceptance	16.8 (7.4)	18.0 (7.8)	18.5 (7.0)	17.2 (7.1)	0.10 (.75)	<0.01 (.96)	0.77 (.38)
CGI	3.4 (1.2)	3.6 (1.1)	3.1 (1.5)	3.0 (1.0)	3.58 (.06)	0.08 (.78)	0.23 (.63)
GAF	63.7 (9.7)	61.8 (10.6)	66.8 (13.5)	65.7 (6.0)	3.0 (.09)	0.54 (.46)	0.03 (.85)

Note. CT = Cognitive Therapy; ACT = Acceptance and Commitment Therapy; BDI = Beck Depression Inventory; BAI = Beck Anxiety Inventory; OQ = Outcome Questionnaire; AAQ = Acceptance and Action Questionnaire; KIMS = Kentucky Inventory of Mindfulness Skills; CGI = Clinical Global Impression; GAF = Global Assessment of Functioning.

and those who did not on key demographic or outcome variables, but those diagnosed with an anxiety disorder were more likely to have completed the follow-up assessment, $\chi^2(3)=8.11$, $p=.04$. A roughly equal percentage of ACT ($n=67\%$) and CT ($n=71\%$) participants completed follow-up, $\chi^2(1)=0.35$, $p=.56$. To probe yet further, a series of analyses were performed in order to determine if the follow-up status (i.e., whether or not a participant was available at follow-up), the treatment condition, or the interaction of follow-up status and condition was related to key demographic, comorbidity, and outcome variables. Logistic regression and ANOVAs indicated that neither follow-up status nor treatment condition nor their interaction was related to baseline characteristics of the sample (Tables 1 and 2).

ANALYTIC STRATEGY

Because we have previously reported posttreatment outcome results on an overlapping sample (Forman et al., 2007), we focus here on follow-up analyses. We conducted long-term follow-up analyses on two samples: (a) an intention-to-treat (ITT) sample ($n=132$) with missing follow-up scores imputed using Expectation Maximization (EM; a relatively sophisticated method of estimating missing values based on iteratively estimating missing data and estimating relationships between observed relationships in the full dataset including, in this case, demographic variables and baseline, posttreatment and observed follow-up values; Dempster, Laird & Rubin, 1977; Schafer & Graham, 2002); and (b) a sample ($n=91$) of those who had presented for follow-up and who were treatment completers (i.e., all those who presented for follow-up were regarded as completers, which we had defined a priori as having received ≥ 5 sessions).⁴

⁴We also considered utilizing Hierarchical Linear Modeling (HLM) because of its ability to model data at the individual and group (i.e., nested) levels and its ability to handle missing data. However, the former was not a focus of the current study and the latter could be even more conservatively handled through the Expectation-Maximization (EM) algorithm (Hedeker & Gibbons, 1997). Of note, we obtained similar results with missing data included and excluded, suggesting that missing data are not having a substantial effect on outcome. Additionally, some experts in HLM caution against its use in naturalistic studies. HLM can lead to a misinterpretation of the rate of progress being made at different points in treatment because it treats participants who end treatment “prematurely” as though they are going forward with therapy after already having made gains when in fact that is not the case (Baldwin, Berkeljon, Atkins, Olsen, & Nielsen, 2009). Finally, in an attempt to be maximally conservative, we re-ran ITT analyses making different assumptions about missing data, and obtained equivalent results (see footnote 5).

LONG-TERM OUTCOMES ACROSS TREATMENTS

Significant and large main effects of time on depression, $F(1, 127)=25.86$, $p<.001$, $\eta_p^2=.16$, $f=.44$; anxiety, $F(1, 127)=16.21$, $p<.001$, $\eta_p^2=.11$, $f=.35$; general functioning, $F(1, 127)=19.55$, $p<.001$, $\eta_p^2=.13$, $f=.39$; and quality of life, $F(1, 124)=4.36$, $p=.02$, $\eta_p^2=.03$, $f=.18$, were obtained from pretreatment to follow-up (and from pre- to posttreatment) in the ITT analyses. The completer analysis revealed similarly sized effects of time on depression, $F(1, 70)=25.56$, $p<.001$, $\eta_p^2=.27$, $f=.61$; anxiety, $F(1, 70)=19.43$, $p<.001$, $\eta_p^2=.22$, $f=.53$; general functioning, $F(1, 70)=19.46$, $p<.001$, $\eta_p^2=.22$, $f=.53$; and quality of life, $F(1, 82)=2.41$, $p=.10$, $\eta_p^2=.03$, $f=.18$. As an ad hoc analysis, diagnostic category (anxiety disorder, depressive disorder, both anxiety and depressive disorders) was entered as a moderator. Results revealed that those diagnosed with an anxiety disorder alone start at higher levels and improve faster on the BAI, $F(1, 68)=3.01$, $p=.01$, $\eta_p^2=.13$, and those diagnosed with comorbid depression and anxiety disorders improve more slowly on the BDI, $F(1, 68)=2.04$, $p=.06$, $\eta_p^2=.09$.

LONG-TERM OUTCOMES BY TREATMENT GROUP

As in our previous analysis, no differences emerged by group at posttreatment. However, at follow-up, a Group \times Time interaction effect emerged. In the full ITT sample, the CT group exhibited greater maintenance of treatment effects for depression, $F(1, 128)=5.36$, $p=.02$, $\eta_p^2=.04$, $f=.21$; general functioning, $F(1, 128)=5.08$, $p=.03$, $\eta_p^2=.04$, $f=.22$; quality of life (weak trend; $F[1, 124]=2.78$, $p=.08$, $\eta_p^2=.02$, $f=.14$); though not for anxiety, $F(1, 128)=.28$, $p=.87$, $\eta_p^2<.01$, $f=.07$). Results for the completers sample followed a similar pattern: general functioning, $F(1, 68)=2.40$, $p=.10$, $\eta_p^2=.03$, $f=.19$; depression, $F(1, 68)=2.75$, $p=.07$, $\eta_p^2=.04$, $f=.21$; quality of life, $F(1, 82)=1.70$, $p=.19$, $\eta_p^2=.02$, $f=.13$; and anxiety, $F(1, 68)=.68$, $p=.51$, $\eta_p^2=.01$, $f=.10$.⁵ Thus, analyses revealed a small- to moderate-sized interaction effect on depression, functioning, and quality of life and a near-zero effect on

⁵Despite the reassuring analyses indicating that follow-up status did not impact outcomes, it is possible that the trajectories of those who presented for follow-up assessments differed from those who did not. To further increase confidence in our findings in light of missing data, we repeated the long-term follow-up analyses using two additional samples derived from alternate assumptions: an ITT sample with baseline values carried forward (which makes the conservative assumption that those who did not present for follow-up regressed to baseline values) and an ITT sample with the last assessment observation carried forward (which makes the assumption that no change occurred between posttreatment and follow-up). Results from these two samples were equivalent to those obtained from the EM and completer samples.

Table 3
Mean Scores and Standard Deviations by Measure

	Baseline		Posttreatment		Follow-up	
	CBT	ACT	CBT	ACT	CBT	ACT
BDI	18.8 (10.2)	19.2 (11.2)	7.6 (8.8)	7.7 (6.8)	7.1 (10.2)	9.2 (9.1)
BAI	15.5 (11.7)	14.6 (9.0)	6.7 (6.6)	7.4 (5.9)	6.1 (5.2)	7.9 (7.3)
OQ	82.1 (17.4)	81.6 (17.1)	66.6 (14.1)	67.1 (13.9)	66.1 (15.6)	68.4 (17.5)
QOLI	0.65 (1.99)	1.12 (2.47)	1.21 (2.35)	1.62 (2.18)	1.60 (1.91)	1.66 (1.91)

Note. BDI = Beck Depression Inventory; BAI = Beck Anxiety Inventory; OQ = Outcome Questionnaire; QOLI = Quality of Life Inventory; CT = Cognitive Therapy; ACT = Acceptance and Commitment Therapy.

anxiety. Observed values at baseline, posttreatment, and follow-up are listed in Table 3. No support was obtained for the hypothesis that either diagnostic status or mindfulness or acceptance measures would further qualify the Time \times Treatment effect ($p_s \geq .16$, $\eta_p^2 \leq .02$), with one exception; KIMS-Describe moderated the Time \times Treatment effect on the QOLI, $F(2, 119) = 13.89$, $p = .003$, $\eta_p^2 = .05$. Completer-only analyses yielded equivalent findings.

CLINICAL SIGNIFICANCE

Clinical significance was analyzed using the recommendations of Jacobson and Truax (1991). We determined the proportion of participants who had “recovered,” i.e., were in the “clinical” range at baseline and “normative” range at follow-up (with cutoff determined by means and standard deviations from normative and clinical samples for each measure; see Jacobson & Truax, 1991) by an amount that exceeded the reliable change index for the measure (determined by reliability and standard deviation of each measure; see Jacobson & Truax). For the BDI, 81.8% of CT patients versus 60.7% of ACT patients reliably recovered. For the BAI, these numbers were 72.7% (CT) and 56.0% (ACT), for OQ they were 46.4% (CT) and 22.6% (ACT), and for QOLI they were 37.8% CT and 22.9% ACT (Figure 2).

Discussion

ACT has demonstrated effectiveness across a wide range of problem behaviors (Hayes, et al., 2006; Powers, et al., 2009); however, ACT intervention studies often lack long-term follow-up analyses and active comparisons. The current study sought to address the question of ACT's long-term effectiveness compared to CT.

The results reveal that the two treatments are equally effective in the short term: both were successful in maintaining improvements in depression, anxiety, and general functioning. Yet, statistical comparisons of long-term outcomes suggest that CT has a slight advantage over ACT in the long-term maintenance of gains, at least with respect to

depressive symptoms and general functioning. By follow-up, quality of life appeared to increase at a steeper rise in those receiving CT (versus ACT), though this result is tempered by the fact that ending points were equivalent. In contrast to some other reports, baseline levels of acceptance and mindfulness did not qualify these effects.

Mean differences in symptom level change were modest (e.g., 3 BDI points); however, clinical significance analyses revealed that roughly one-third more CT than ACT patients remained in the recovered range at long-term follow-up, with similar differences for quality of life. Furthermore, in the case of the OQ—which is a good all-around indicator of interpersonal and occupational functioning as well as symptom distress—more than twice as many CT as ACT patients were in the normative range at follow-up.

Several possible explanations can be offered for these findings. First, it is possible that CT is a more intuitive and simpler intervention, thus facilitating the ability of patients to apply the learned strategies and skills independently once treatment has terminated. As we have argued elsewhere (Forman & Herbert, 2009), CT is highly compatible with folk psychology,




FIGURE 2 Percent of participants (in clinical range at baseline) who have reliably recovered at follow-up. Note. BDI = Beck Depression Inventory; BAI = Beck Anxiety Inventory; OQ = Outcome Questionnaire; QOLI = Quality of Life Inventory; CT = Cognitive Therapy; ACT = Acceptance and Commitment Therapy.

whereas ACT tends to be more counterintuitive. In a sense, ACT requires the individual to apply concepts that often run counter to prevailing social norms and customs, and as the time away from the therapist's support and guidance increases, the more difficult it may be for some individuals to implement ACT principles in the face of these prevailing contingencies. Another possibility is that study therapists were better able to administer CT (although the comparable competence and adherence data do not support this hypothesis), perhaps because they were relatively inexperienced (though well-trained). Finally, most of the patients in this study were relatively high functioning and had only modest symptom levels. Some authors have speculated that acceptance-based psychotherapies are especially well suited to more severe, treatment-resistant patients (Kenny & Williams, 2007; Twohig, 2009). No matter the explanation, it challenges the notion that ACT contains more active ingredients than does Beckian CT or that its principles have strong empirical backing.

The present study is the first to our knowledge to directly compare the long-term efficacy of ACT to another gold-standard CBT. Strengths of the study include true random assignment, the long-term follow-up assessment, and therapists whose adherence, competence, and absence of treatment allegiance have been ensured. The study also emphasized external validity using minimal exclusion criteria, and a naturalistic course of treatment based on general guidelines instead of highly prescriptive manuals. In addition, results were verified across multiple imputed datasets with varying assumptions. These strengths are tempered by complementary limitations including diagnostic variability, intervention variability due to minimalist treatment manualization, relatively novice therapists, the lack of a no-treatment control and the lack of formal mediational analyses. The fact that most measures were self-report represents an additional weakness in that data can be limited by self-knowledge and/or response bias (and can reflect experimenter demand). The absence of data on subsequent treatment during the follow-up period should also be noted. In addition, retention was modest (though comparable to similar trials), likely because of the high mobility of the sample. The convergence of findings across various ITT and completer samples partially attenuates this concern.

The results of this study raise interesting questions about the long-term maintenance of gains in a newer, acceptance-based model of CBT relative to standard CT. These findings merit further study, and replication is needed before definitive conclusions can be drawn. Such research would be augmented by an even longer follow-up period and blind clinical

assessments, as well as a more clinically impaired sample treated by expert therapists. Studies of moderating variables are especially needed to assess the potential advantages of matching patient characteristics with the intervention most likely to yield long-term benefits.

References

- Arch, J. J., & Craske, M. G. (2008). Acceptance and commitment therapy and cognitive behavioral therapy for anxiety disorders: Different treatments, similar mechanisms? *Clinical Psychology: Science and Practice*, 5, 263–279.
- Bach, P., & Hayes, S. C. (2002). The use of acceptance and commitment therapy to prevent the rehospitalization of psychotic patients: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 70(5), 1129–1139.
- Baer, R. A., Smith, G. T., & Allen, K. B. (2004). Assessment of mindfulness by self-report: The Kentucky inventory of mindfulness skills. *Assessment*, 11, 191–206. <http://dx.doi.org/10.1177/1073191104268029>
- Baldwin, S. A., Berkeljon, A., Atkins, D. C., Olsen, J. A., & Nielsen, S. L. (2009). Rates of change in naturalistic psychotherapy: Contrasting dose-effect and good-enough level models of change. *Journal of Consulting and Clinical Psychology*, 77, 203–211. <http://dx.doi.org/10.1037/A0015235>
- Beck, A. T. (1991). Cognitive therapy: A 30 year retrospective. *American Psychologist*, 46(4), 368–375.
- Beck, A. T., Epstein, N., Brown, G. K., & Steer, R. A. (1988). An inventory for measuring anxiety: Psychometric properties. *Journal of Consulting & Clinical Psychology*, 56, 893–897.
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: Guilford Press.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Beck, A. T., Steer, R. A., & Garbin, M. G. (1988). Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*, 8(1), 77–100.
- Blackledge, J. T., & Hayes, S. C. (2006). Using acceptance and commitment training in the support of parents of children diagnosed with autism. *Child & Family Behavior Therapy*, 28(1), 1–18.
- Block, J. A. (2003). *Acceptance or change of private experiences: A comparative analysis in college students with public speaking anxiety*. Unpublished doctoral dissertation, State University of New York at Albany.
- Brown, L. A., Forman, E. M., Herbert, J. D., Hoffman, K. L., Yuen, E. K., & Goetter, E. M. (2011). A randomized controlled trial of acceptance-based behavior therapy and cognitive therapy for test anxiety: A pilot study. *Behavior Modification*, 35, 31–53. <http://dx.doi.org/10.1177/0145445510390930>
- Brown, R. A., Lejuez, C., Kahler, C. W., Strong, D. R., & Zvolensky, M. J. (2005). Distress tolerance and early smoking lapse. [Original]. *Clinical Psychology Review*, 25(6), 713–733.
- Butler, A. C., Chapman, J. E., Forman, E. M., & Beck, A. T. (2006). The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*, 26(1), 17–31.
- Corrigan, P. W. (2001). Getting ahead of the data: A threat to some behavior therapies. *the Behavior Therapist*, 24(9), 189–193.
- Corrigan, P. W. (2002). The data is still the thing: A reply to Gaynor and Hayes. *the Behavior Therapist*, 25, 140.

- Dahl, J., Wilson, K. G., & Nilsson, A. (2004). Acceptance and commitment therapy and the treatment of persons at risk for long-term disability resulting from stress and pain symptoms: A preliminary randomized trial. *Behavior Therapy, 35*(4), 785–801.
- Dalrymple, K. L., & Herbert, J. D. (2007). Acceptance and commitment therapy for generalized social anxiety disorder: A pilot study. *Behavior Modification, 31*(5), 543–568.
- Dempster, A. P., Laird, N. M., & Rubin, D. B. (2004). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society, B, 39*, 1–38.
- Dimidjian, S., Hollon, S. D., Dobson, K. S., Schmalzing, K. B., Kohlenberg, R. J., Addis, M. E., . . . Jacobson, N. S. (2006). Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *Journal of Consulting and Clinical Psychology, 74*, 658–670. <http://dx.doi.org/10.1037/0022-006x.74.4.658>
- Durham, R. C., Higgins, C., Chambers, J. A., Swan, J. S., & Dow, M. G. (2011). Long-term outcome of eight clinical trials of CBT for anxiety disorders: Symptom profile of sustained recovery and treatment-resistant groups. *Journal of Affective Disorders. <http://dx.doi.org/10.1016/j.jad.2011.09.017>*
- First, M. B., Williams, J. B. W., Spitzer, R. L., & Gibbon, M. (2007). *Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Clinical Trials Version (SCID-CT)*. New York: Biometrics Research.
- Forman, E. M., Butryn, M. L., Hoffman, K. L., & Herbert, J. D. (2009). An open trial of an acceptance-based behavioral intervention for weight loss. *Cognitive and Behavioral Practice, 16*(2), 223–235.
- Forman, E. M., & Herbert, J. D. (2009). New directions in cognitive behavior therapy: Acceptance-based therapies. In W. T. O'Donohue, & J. E. Fisher (Eds.), *General principles and empirically supported techniques of cognitive behavior therapy* (pp. 77–101). Hoboken, NJ: Wiley & Sons.
- Forman, E. M., Herbert, J. D., Moitra, E., Yeomans, P. D., & Geller, P. A. (2007). A randomized controlled effectiveness trial of acceptance and commitment therapy and cognitive therapy for anxiety and depression. *Behavior Modification, 31*, 772–799. <http://dx.doi.org/10.1177/0145445507302202>
- Frisch, M. B., Clark, M. P., Rouse, S. V., Rudd, M. D., Pawelek, J. K., Greenstone, A., & Kopplin, D. A. (2005). Predictive and treatment validity of life satisfaction and the quality of life inventory. *Assessment, 12*, 66–78. <http://dx.doi.org/10.1177/1073191104268006>
- Gaudiano, B. A. (2009). Öst's (2008) methodological comparison of clinical trials of acceptance and commitment therapy versus cognitive behavior therapy: Matching apples with oranges? *Behaviour Research and Therapy, 47*, 1066–1070.
- Gaudiano, B. A. (2009b). Reinventing the wheel versus avoiding past mistakes when evaluating psychotherapy outcome research: Rejoinder to Öst (2009). *Unpublished manuscript*.
- Gaudiano, B. A., & Herbert, J. D. (2006). Acute treatment of inpatients with psychotic symptoms using Acceptance and Commitment Therapy: Pilot results. *Behaviour Research and Therapy, 44*, 415–437. <http://dx.doi.org/10.1016/j.brat.2005.02.007>
- Gifford, E. V., Kohlenberg, B. S., Hayes, S. C., Antonuccio, D. O., Piasecki, M. M., Rasmussen-Hall, M. L., & Palm, K. M. (2004). Acceptance-based treatment for smoking cessation. *Behavior Therapy, 35*(4), 689–705.
- Gillis, M. M., Haaga, D. A., & Ford, G. T. (1995). Normative values for the Beck Anxiety Inventory, Fear Questionnaire, Penn State Worry Questionnaire, and Social Phobia and Anxiety Inventory. *Psychological Assessment, 7*, 450–455. <http://dx.doi.org/10.1037/1040-3590.7.4.450>
- Gloaguen, V., Cottraux, J., Cucherat, M., & Blackburn, I. M. (1998). A meta-analysis of the effects of cognitive therapy in depressed patients. *Journal of Affective Disorders, 49*(1), 59–72.
- Guy, W. (1976). *ECDEU Assessment Manual for Psychopharmacology*. Rockville, MD: National Institute of Mental Health.
- Hayes, S. C. (2002). On being visited by the vita police: A reply to Corrigan. *the Behavior Therapist, 25*, 134–137.
- Hayes, S. C. (2008). Climbing our hills: A beginning conversation on the comparison of acceptance and commitment therapy and traditional cognitive behavioral therapy. *Clinical Psychology: Science and Practice, 15*(4), 286–295.
- Hayes, S. C., Levin, M., Plumb, J., Boulanger, J., & Pistorello, J. (in press). Acceptance and commitment therapy and contextual behavioral science: Examining the progress of a distinctive model of behavioral and cognitive therapy. *Behavior Therapy. <http://dx.doi.org/10.1016/j.beth.2009.08.002>*
- Hayes, S. C., Luoma, J. B., Bond, F. W., Masuda, A., & Lillis, J. (2006). Acceptance and commitment therapy: Model, processes and outcomes. *Behavior Research and Therapy, 44*(1), 1–25.
- Hayes, S. C., Strosahl, K., & Wilson, K. G. (1999). *Acceptance and commitment therapy: An experiential approach to behavior change*. New York: Guilford Press.
- Hayes, S. C., Strosahl, K., Wilson, K. G., Bissett, R. T., Pistorello, J., Toarmino, D., . . . McCurry, S. M. (2004). Measuring experiential avoidance: A preliminary test of a working model. *Psychological Record, 54*(4), 553–578.
- Hayes, S. C., Villatte, M., Levin, M., & Hildebrandt, M. (2011). Open, aware, and active: Contextual approaches as an emerging trend in the behavioral and cognitive therapies. *Annual Review of Clinical Psychology, 7*, 141–168. <http://dx.doi.org/10.1146/annurev-clinpsy-032210-104449>
- Hayes, S. C., Wilson, K. G., Gifford, E. V., Bissett, R. T., Piasecki, M., Batten, S. V., . . . Gregg, J. (2004). A preliminary trial of twelve-step facilitation and acceptance and commitment therapy with polysubstance-abusing methadone-maintained opiate addicts. *Behavior Therapy, 35*, 667–688. <http://dx.doi.org/10.1016/S0005-7894%2804%2980014-5>
- Hedeker, D., & Gibbons, R. D. (1997). Application of random-effects pattern-mixture models for missing data in longitudinal studies. *Psychological Methods, 2*(1), 64–78.
- Herbert, J. D., & Forman, E. M. (Eds.). (2011). *Acceptance and mindfulness in cognitive behavior therapy*. Hoboken, NJ: Wiley.
- Herbert, J. D., & Forman, E. M. (in press). Caution: The differences between CT and ACT may be larger (and smaller) than they appear. *Behavior Therapy. <http://dx.doi.org/10.1016/j.beth.2009.09.005>*
- Hernández-López, M., Luciano Soriano, M. C., Bricker, J. B., Roales-Nieto, J. G., & Montesinos Marin, F. (2009). Acceptance and commitment therapy for smoking cessation: A preliminary study of its effectiveness in comparison with cognitive behavioral therapy. *Psychology of Addictive Behaviors, 23*(4), 723–730.
- Hernandez Lopez, M., Roales Nieto, J. G., Luciano Soriano, M. C., & Montesinos Marin, F. (2005). A comparison between two psychological treatments oriented to smoking-cessation: Acceptance versus control strategies. *Paper presented at the Annual Convention of the Association for Behavior Analysis, Chicago, IL*.
- Hofmann, S. G., & Asmundson, G. (2008). Acceptance mindfulness-based therapy: New wave or old hat? *Clinical Psychology Review, 28*, 1–16.
- Hofmann, S. G., & Asmundson, G. J. (in press). The science of cognitive-behavioral therapy. *Behavior Therapy*.

- Hofmann, S. G., & Smits, J. A. J. (2008). Cognitive-behavioral therapy for adult anxiety disorders: A meta-analysis of randomized placebo-controlled trials. *Journal of Clinical Psychiatry*, *69*(4), 621–632.
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, *59*(1), 12–19.
- Kabat-Zinn, J. (1990). *Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness*. New York: Delacourt.
- Kenny, M. A., & Williams, J. M. G. (2007). Treatment-resistant depressed patients show a good response to mindfulness-based cognitive therapy. *Behaviour Research and Therapy*, *45*, 617–625. <http://dx.doi.org/10.1016/j.brat.2006.04.008>
- Lambert, M. J., Burlingame, G. M., Umphress, V., Hansen, N. B., Vermeersch, D. A., Clouse, G. C., & Yanchar, S. C. (1996). The reliability and validity of the outcome questionnaire. *Clinical Psychology & Psychotherapy*, *3*(4), 249–258.
- Lambert, M. J., Hansen, N., Umphress, V., Lunnen, K., Okiishi, J., Burlingame, G., . . . Reisinger, C. (1996). *Administration and scoring manual for the Outcome Questionnaire (OQ 45.2)*. Wilmington, DE: American Professional Credentialing Services.
- Lappalainen, R., Lehtonen, T., Skarp, E., Taubert, E., Ojanen, M., & Hayes, S. C. (2007). The impact of CBT and ACT models using psychology trainee therapists: a preliminary controlled effectiveness trial. *Behavior Modification*, *31*, 488–511.
- Lecrubier, Y., Sheehan, D. V., Weiller, E., Amorim, P., Bonora, I., Sheehan, K. H., . . . Dunbar, G. C. (1997). The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: Reliability and validity according to the CIDI. *European Psychiatry*, *12*(5), 224–231.
- Lillis, J., Hayes, S. C., Bunting, K., & Masuda, A. (2009). Teaching acceptance and mindfulness to improve the lives of the obese: A preliminary test of a theoretical model. *Annals of Behavioral Medicine*, *37*, 58–69.
- Linehan, M. M. (1993). *Cognitive-behavioral treatment of borderline personality disorder*. New York, NY: Guilford Press.
- Longmore, R. J., & Worrell, M. (2007). Do we need to challenge thoughts in cognitive behavior therapy? *Clinical Psychology Review*, *27*, 173–187. <http://dx.doi.org/10.1016/j.cpr.2006.08.001>
- McCracken, L. M., MacKichan, F., & Eccleston, C. (2007). Contextual cognitive-behavioral therapy for severely disabled chronic pain sufferers. *European Journal of Pain*, *11*(3), 314–322.
- McGrath, K. B., Forman, E. M., & Herbert, J. D. (2009, November). Validation of the ACT/CT adherence and competence rating scale: Phase two. *Paper presented at the 43rd Annual Convention of the Association for Behavioral and Cognitive Therapies, New York, NY*.
- Norcross, J. C., Hedges, M., & Castle, P. H. (2002). Psychologists conducting psychotherapy in 2001: A study of the Division 29 membership. *Psychotherapy: Theory, Research, Practice, Training*, *39*(1), 97–102.
- Ossman, W. A., Wilson, K. G., Storaasli, R. D., & McNeill, J. W. (2006). A preliminary investigation of the use of acceptance and commitment therapy in group treatment for social phobia. *International Journal of Psychology and Psychological Therapy*, *6*(3), 397–416.
- Öst, L. (2008). Efficacy of the third wave of behavioral therapies: A systematic review and meta-analysis. *Behaviour Research and Therapy*, *46*, 296–321. <http://dx.doi.org/10.1016/j.brat.2007.12.005>
- Öst, L. (2009). Inventing the wheel once more or learning from the history of psychotherapy research methodology: Reply to Gaudiano's comments on Öst's (2008) review. *Behaviour Research and Therapy*, *47*, 1071–1073.
- Powers, M. B., Zum Vörde Sive Vörding, M. B., & Emmelkamp, P. (2009). Acceptance and commitment therapy: A meta-analytic review. *Psychotherapy and Psychosomatics*, *78*(2), 73–80.
- Roemer, L., & Orsillo, S. M. (2005). An acceptance based behavior therapy for generalized anxiety disorder. In S. M. Orsillo, & L. Roemer (Eds.), *Acceptance and mindfulness-based approaches to anxiety: Conceptualization and treatment*. New York: Springer.
- Roemer, L., Salters-Pedneault, K., & Orsillo, S. M. (2006). Incorporating mindfulness- and acceptance-based strategies in the treatment of generalized anxiety disorder. In Baer, R. A. (Ed.), *Mindfulness-based treatment approaches: Clinician's guide to evidence base and applications* (pp. 51–74). San Diego, CA: Elsevier Academic Press
- Schafer, J. L., & Graham, J. W. (2002). Missing data: Our view of the state of the art. *Psychological Methods*, *7*(2), 147–177.
- Segal, D. L., Coolidge, F. L., Cahill, B. S., & O'Riley, A. A. (2008). Psychometric properties of the Beck Depression Inventory-II (BDI-II) among community-dwelling older adults. *Behavior Modification*, *32*, 3–20. <http://dx.doi.org/10.1177/0145445507303833>
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guilford Press.
- Shapiro, D. A., Barkham, M., Rees, A., Hardy, G. E., Reynolds, S., & Startup, M. (1994). Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic interpersonal psychotherapy. *Journal of Consulting and Clinical Psychology*, *62*(3), 522–534.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., . . . Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (MINI): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, *59*, 22–33.
- Spitzer, R. L., Gibbon, M., Williams, J. B. W., & Endicott, J. (1996). Global assessment of functioning (GAF) scale. In L. I. Sederer, & B. Dickey (Eds.), *Outcomes assessment in clinical practice* (pp. 76–78). Baltimore, MD: Williams and Wilkins.
- Stotts, A. L., Masuda, A., & Wilson, K. G. (2009). Using acceptance and commitment therapy during methadone dose reduction: Rational, treatment description, and a case report. *Cognitive and Behavioral Practice*, *16*(2), 205–213.
- Tapper, K., Shaw, C., Ilsley, J., Hill, A. J., Bond, F. W., & Moore, L. (2009). Exploratory randomised controlled trial of a mindfulness-based weight loss intervention for women. *Appetite*, *52*(2), 396–404.
- Twohig, M. P. (2008). A randomized clinical trial of acceptance and commitment therapy versus progressive relaxation training in the treatment of obsessive compulsive disorder. *Dissertation Abstracts International: Section B: The Sciences & Engineering*, *68*(2008), 4850.
- Twohig, M. P. (2009). Acceptance and commitment therapy for treatment-resistant posttraumatic stress disorder: A case study. *Cognitive and Behavioral Practice*, *16*(3), 243–252.
- Twohig, M. P., Hayes, S. C., & Masuda, A. (2006). Increasing willingness to experience obsessions: acceptance and commitment therapy as a treatment for obsessive-compulsive disorder. *Behavior Therapy*, *37*(1), 3–13.
- Twohig, M. P., Shoenberger, D., & Hayes, S. C. (2007). A preliminary investigation of acceptance and commitment therapy as a treatment for marijuana dependence in adults. *Journal of Applied Behavior Analysis*, *40*(4), 619–632.

- Vowles, K. E., & McCracken, L. M. (2008). Acceptance and values-based action in chronic pain: A study of treatment effectiveness and process. *Journal of Consulting & Clinical Psychology, 76*(3), 397–407.
- Whisman, M. A., Perez, J. E., & Ramel, W. (2000). Factor structure of the Beck Depression Inventory—Second Edition (BDI-II) in a student sample. *Journal of Clinical Psychology, 56*(4), 545–551. <http://dx.doi.org/10.1002/%28SICI%291097-4679%28200004%2956:4%3C545::AID-JCLP7%3E3.0.CO;2-U>
- Woods, D. W., Wetterneck, C. T., & Flessner, C. A. (2006). A controlled evaluation of acceptance and commitment therapy plus habit reversal for trichotillomania. *Behavior Research and Therapy, 44*(5), 639–656.
- Zettle, R. D. (2003). Acceptance and commitment therapy (ACT) vs. systematic desensitization in treatment of mathematics anxiety. *Psychological Record, 53*(2), 197–215.
- Zettle, R. D., & Hayes, S. C. (1986). Dysfunctional control by client verbal behavior: The context of reason giving. *The Analysis of Verbal Behavior, 4*, 30–38.

RECEIVED: July 9, 2011

ACCEPTED: April 16, 2012

Available online 24 April 2012